

**1) Emergency Symptoms/signs**

Thunderclap onset  
Accelerated/Malignant hypertension  
Acute onset with papilloedema  
Acute onset with focal neurological signs  
Head trauma with raised ICP headache  
Photophobia + nuchal rigidity + fever +/-rash  
Reduced consciousness  
Acute red eye: ?acute angle closure glaucoma

**New** onset headache in:

- 3rd trimester pregnancy/early postpartum
- Significant head injury – especially elderly patients, alcohol dependency, people on anticoagulants

**2) Giant Cell arteritis**

Incidence 2/10,000 per year  
• Consider with presentations of new headache in >50 year olds  
• Many headaches respond to high dose steroids **NB** do not use response as the sole diagnostic factor.  
• ESR can be normal in 10% - check CRP as well  
• Symptoms may include: jaw/tongue claudication, visual disturbance, temporal artery: prominent, tender, diminished pulse; other cranial nerve palsies, limb claudication

**Urgent referral to:**

- Rheumatology if diagnosis clear
- Neurology if headache or possibly GCA
- Ophthalmology if amaurosis fugax / visual loss / diplopia **NOT** migrainous auras

**3) 2WW - suspected cancer referral**

• **Headache with features of raised intracranial pressure:-**

- Actively wakes a patient from sleep, but not migraine or cluster
- **Precipitated** by Valsalva manoeuvres i.e. cough, straining at stool
- Papilloedema
- Other symptoms of raised ICP headache including
  - Headache present upon waking and easing once up (analgesic overuse can cause this pattern) and worse when recumbent
  - Pulse synchronous tinnitus
  - Episodes of transient visual loss when changing posture e.g. upon standing
  - Vomiting - significance should be judged in context as nausea and vomiting are features of migraine

• **Headache with new onset seizures**

• **Headache with persistent new or progressive neurological deficit**

• **A relevant history of malignancy which might have metastasised to the brain**

• **Vomiting without other obvious cause (i.e. not just due to migraine)**

**4) Red Flags (for secondary headaches)**

- Headache rapidly increasing in severity and frequency despite appropriate treatment
- Undifferentiated headache (not migraine / tension headache) of recent origin and present for >8 weeks
- Recurrent headaches triggered by exertion
- Orthostatic Headache (headache that occurs in the upright position, suggesting low CSF pressure)
- New onset headache in:-
  - >50 years old (consider giant cell arteritis)
  - Immunosuppressed / HIV

**5) Migraine**

- Throbbing pain lasting hours - 3 days
  - Sensitivity to stimuli: light and sound, sometimes smells
  - Nausea
  - Aggravated by physical activity (prefers to lie/sit still)
- Aura, if present, that evolves slowly (in contrast to TIA/stroke) and lasts minutes - 60min

**'Chronic Migraine'**

≥15 headache days/month of which ≥8 are migraine

**Acute treatments:**

Aspirin dispersible 900mg or NSAID, taken with metoclopramide or domperidone **NB** Note MHRA warning

[MHRA \(2014\): Domperidone: risks of cardiac side effects](#)

[MHRA \(2013\) Metoclopramide: risk of neurological adverse effects](#)

A triptan but <10 days per month (best <6/month)

Don't use opiates as they tend to lead to increase nausea and lead to an overuse headache

**Tension Type Headache**

Band-like ache, mostly featureless  
Can have mild photo OR phonophobia but NO nausea  
Many believe this is simply a milder form of migraine i.e. same biology and thus similar treatments can be effective

**Cluster Headache**

More common in men  
Most severe pain ever lasting 30-120 minutes  
Unilateral, side-locked  
Agitation, pacing **NB** migraineurs prefer to keep still  
Unilateral Cranial Autonomic features:-  
tearing, red conjunctiva, ptosis, miosis, nasal stuffiness

**Acute treatments:**

Sumatriptan injection 6mg s.c. - contra-indicated for IHD and stroke

Hi-flow oxygen through a non-rebreathe bag and mask

Prednisolone 60mg od for 1 week can abort a bout of attacks

**Triptan Overuse Headache**

Can be migrainous and/or tension type  
Triptan intake: ≥10 days/month for ≥3 months

**Treatment:** Stop triptan for 2-3 months

**Analgesic Overuse Headache**

Can be migrainous and/or tension type  
Analgesic intake ≥15 days/month (opiates ≥10 days)

For ≥3 consecutive months

**Treatment:** stop analgesic for 3 months

**6) Botulinum Toxin for Chronic Migraine: (NICE TA260)**

Between 31 and 39 injections i.m. around scalp and neck every 12 weeks

Minimum treatment criteria:

- Chronic migraine i.e. ≥15 headache days/month of which ≥8 are migraine for a minimum of 3 consecutive months
- Tried 3 different migraine preventatives at maximally tolerated doses for 3 months each **not** including pizotifen
- Not overusing triptans, opiates or other analgesics